

Extended supersolid phase of frustrated hard-core bosons on a triangular lattice

Fa Wang, Frank Pollmann, Ashvin Vishwanath¹

¹Department of Physics, University of California, Berkeley, CA94720

(Dated: September 9, 2008)

We study a model of hard-core bosons with *frustrated* nearest-neighbor hopping (t) and repulsion (V) on the triangular lattice. We argue for a supersolid ground state in the large repulsion ($V \gg |t|$) limit where a dimer representation applies, by constructing a unitary mapping to the well understood unfrustrated hopping case. This generalized 'Marshall sign rule' allows us to establish the precise nature of the supersolid order by utilizing a recently proposed dimer variational wavefunction, whose correlations can be efficiently calculated using the Grassman approach. By continuity, a supersolid is predicted over the wide parameter range, $V > -2t > 0$. This also establishes a simple phase diagram for the triangular lattice spin 1/2 XXZ antiferromagnet.

PACS numbers:

Supersolidity, where superfluid and crystalline orders coexist, have fascinated physicists since they were first theoretically proposed [1]. Recent experimental results in ⁴He [2] that are still under active debate have led to renewed interest. Experimental developments on a different front, in the realization of optical lattices in ultracold atomic systems, motivated a search for a lattice supersolid. One of the more promising candidates is a model of strongly interaction hard-core bosons on a triangular lattice. The model Hamiltonian reads

$$H = -t \sum_{\langle ij \rangle} (b_i^\dagger b_j + \text{H.c.}) + V \sum_{\langle ij \rangle} \left(n_i - \frac{1}{2} \right) \left(n_j - \frac{1}{2} \right), \quad (1)$$

where b_i ($b_i^\dagger$) annihilates (creates) a hard-core boson on site i and $n_i = b_i^\dagger b_i$ are density operators. The model is equivalent to the XXZ spin-1/2 Hamiltonian on the triangular lattice

$$H = \sum_{\langle ij \rangle} \left[\frac{J_\perp}{2} (s_i^+ s_j^- + s_i^- s_j^+) + \frac{J_z}{4} s_i^z s_j^z \right], \quad (2)$$

where $s^\pm = (1/2)(s^x \pm i s^y)$, $s^{x,y,z}$ are the Pauli matrices and are related to bosons by $s_i^z = (2n_i - 1)/2$, $s_i^+ = b_i^\dagger$, $s_i^- = b_i$, and $J_z = V$, $J_\perp = -2t$. The discussion below will be largely in terms of the bosons, although we will sometimes switch to the equivalent spin description, when that is more natural. The $t > 0$ case corresponds to the unfrustrated hard-core boson model with repulsive nearest-neighbor interactions. For this case, a variety of studies including large scale quantum Monte Carlo simulations [3, 4, 5] indicate a supersolid phase for all $V/t \geq 8.9$, stabilized by an 'order by disorder' mechanism. The solid order is of the three sublattice ($++-$) type, where two sublattices have the same boson density.

The case of *frustrated* hopping ($t < 0$) suffers from a sign problem in the occupation number basis, and its ground state has been a subject of conjecture for the last three decades. The corresponding spin model is just the XXZ antiferromagnet, which, in the large J_z limit was at the center of the RVB spin liquid proposal of Fazekas and Anderson [6]. Later semiclassical and small cluster numerical studies suggested magnetic order [7], and general arguments which apply to the phase structure of bipartite dimer models, to which this model

can be mapped in the large J_z limit, also indicate the same result [8, 9]. However, the precise nature of ordering has not been conclusively established. In this letter, we show how this problem can be tackled, which is summarized briefly in boson language below. Due to frustration, the ground states in the $V \rightarrow \infty$ limit is extensively degenerate. Within this ground state manifold, we demonstrate that the frustrated problem with $t < 0$ can be mapped, via a nontrivial unitary transformation, onto the unfrustrated one with $t > 0$. Since the latter is well understood [3, 4, 5, 10], many properties of the frustrated case can be immediately derived. Such a generalized 'Marshall sign' was conjectured earlier based on state enumeration and numerics [6, 11]. Here we construct the explicit transformation which proves this conjecture, and moreover utilize it to deduce properties of the frustrated model. Our unitary transformation is diagonal in the occupation number basis, which, combined with our knowledge of the unfrustrated model, allows us to argue that a supersolid state is realized for $t < 0$ as well. The precise details of the superfluid phase ordering requires further calculation. This is carried out using a variational wave-function approach [12] recently introduced for the unfrustrated $t/V = 0^+$ limit, which captures the essential aspects of supersolid order very well and has good variational energy as compared to the quantum Monte Carlo results. Applying the unitary transformation, we obtain a variation wave-function for the frustrated problem. Properties of this wave-function, in particular the phase correlations, are then calculated. The state is found to be a supersolid and the resulting structure of the long range order (LRO) is shown in FIG. 4a. Surprisingly, the superfluid amplitude vanishes on one of the sublattices and hence superfluidity lives exclusively on the honeycomb lattice formed by the remaining two sublattices, on which the amplitude alternates in sign. Contrary to naive expectations, the superfluid amplitude on these sites *exceeds* the maximum superfluid amplitude of the unfrustrated case.

Finally, with this information in hand, we propose a phase diagram for the entire $t/V > 0$ parameter range. Note the point $t/V = 1/2$ corresponds to the spin-isotropic triangular antiferromagnet, where the 120° state is established. This can be smoothly connected to the large V supersolid state derived here as shown in FIG. 1. Supersolid order would




FIG. 1: Schematic phase diagram for both unfrustrated ($t > 0$) and frustrated hopping ($t < 0$) with repulsive interactions ($V > 0$). The three arrows are order parameters $\vec{s} = (b^\dagger + b, ib - ib^\dagger, 2n - 1)$ on the three sub-lattices. For $t/V < -1/2$ or $t/V > 0.1$ there is only superfluid LRO (XY spin order). The thick line $-1/2 < t/V < 0.1$ is the region of supersolid order. $t/V = -1/2$ is the SU(2) symmetric antiferromagnet.

then naturally be preserved over the wide parameter range $0 < -t < V/2$, in contrast to the unfrustrated case, where it is only present for $t < V/10$. The frustrated triangular lattice boson model therefore appears to be an appealing candidate for the realization of the elusive supersolid phase - experimental prospects are discussed at the end. Note, this is also a phase diagram for the spin 1/2 XXZ magnet, and the regime of proposed RVB phase of Fazekas-Anderson [6] is actually a particular spin ordered state.

Strong Repulsion Limit and Generalized Marshall Sign: In the limit of $V \gg |t|$, we can restrict the Hilbert space to a manifold of states which correspond to classical Ising ground states of the triangular antiferromagnet [13]. Every such Ising configurations $\mathcal{S}$ can be represented by a close-packed dimer configuration $\mathcal{C}$ on the dual honeycomb lattice. This is a two-to-one mapping because of the Ising $\mathbb{Z}_2$ symmetry (particle-hole symmetry in the boson language). The Hamiltonian (2), projected into this degenerate subspace, introduces dynamics which splits the degeneracy. Note, to first order in degenerate perturbation theory, only the hopping term $H_t = -t \sum_{\langle ij \rangle} (b_i^\dagger b_j + H.c.)$ plays a role, leading to the double-hexagon resonance in FIG. 2 (a) with amplitude $-t$. The problem of the large repulsion limit is therefore related to finding the ground state of a quantum dimer model with such dimer resonances. We have already noted that the $t > 0$ case is tractable by Quantum Monte Carlo methods since there is no sign problem. However, the problem of interest here is the case $t < 0$. If there is a unitary transformation which changes the sign of every matrix element of H_t , the problem can be mapped to unfrustrated case. This is generically not possible but, *within* the restricted Hilbert space, this indeed happens, and the required unitary transformation is the following. Consider the lattice in FIG. 2 (c) with 1/4 special edges marked as thick and green. One can check by inspection that any double-hexagon resonance will change the number of covered special edges by ± 2 . Therefore, if we define a unitary transformation on the dimer basis

$$|\mathcal{C}'\rangle = U|\mathcal{C}\rangle = i^{N_s(\mathcal{C})}|\mathcal{C}\rangle, \quad (3)$$

where $i = \sqrt{-1}$, and $N_s(\mathcal{C})$ is the number of special (green) edges covered by a dimer in the dimer configuration $\mathcal{C}$, the




FIG. 2: (Color online) (a): two double-hexagon resonance configurations c_{ij} and $\bar{c}_{ij} = c_{ji}$. Red thick bars denote dimers. (b): Kasteleyn orientation and edge weights of the honeycomb lattice. Thick blue edges have weight z , others have weight 1. The green dash-line rhombus encloses the enlarged unit cell. x, y are the principal axis. We use the six sites on a thick-edge hexagon as the basis, labeled as $1, \dots, 6$ as shown in the right-bottom corner. (c): special edges (thick green on the honeycomb) for the unitary transformation relating the unfrustrated and frustrated case. Thin solid green bonds on the triangular lattice are dual to the special edges.

sign of the Hamiltonian will be changed. The unitary transformation does not change the energy spectrum nor correlations that are diagonal in boson density. Hence, thermodynamics - that only depends on energy eigenvalues- is unchanged, for eg. transition temperature and nature of transitions. However, off-diagonal correlations are affected. We can therefore immediately conclude that the ground state has the same three sublattice density modulation as the supersolid phase in the unfrustrated model. Moreover, it also has a finite compressibility, identical to that in the unfrustrated problem, since this can also be expressed as a density-density correlation function. The latter strongly suggests superfluid long range order (a 2D bosonic phase with finite compressibility at zero temperature), and taken all together this points towards supersolid order for $t/V = 0^-$ as well. In order to directly establish off diagonal long range order, and obtain more detailed quantitative information, we turn to a variational wavefunction approach.

Variational Wavefunction We denote the two Ising states related to the dimer state $\mathcal{C}$ as $\mathcal{S}[\mathcal{C}]$ and $\bar{\mathcal{S}}[\mathcal{C}]$ and consider the following kind of wavefunctions,

$$|\Psi\rangle = \sum_{\mathcal{C}} \phi(\mathcal{C})|\mathcal{C}\rangle = \sum_{\mathcal{C}} \phi(\mathcal{C}) \cdot (|\mathcal{S}[\mathcal{C}]\rangle + |\bar{\mathcal{S}}[\mathcal{C}]\rangle) / \sqrt{2} \quad (4)$$

where $\phi(\mathcal{C})$ is the (complex) amplitude. In the dimer representation, the projected H_t corresponds to the double-hexagon resonance in FIG. 2 (a). Only those dimer configurations with 'resonatable' double hexagons appear in the Hamiltonian matrix elements. We denote by c_{ij} , that a particular dimer covering has a resonatable double hexagon at the pair of adjacent plaquettes i, j , where i is the plaquette with two dimers. Under resonance $c_{ij} \rightarrow \bar{c}_{ij} = c_{ji}$. However, the rest of the dimer configuration with this pair of plaquettes removed d_{ij} remains unchanged. Hence, the entire dimer configuration may be denoted as $c_{ij} + d_{ij}$. Note, a single dimer configuration may

have many representations in this notation - one for each resonatable hexagon pair. The variational energy $E = \langle \Psi | H_t | \Psi \rangle$ is

$$E = -t \sum_{\langle ij \rangle} \sum_{d_{ij}} [\phi^*(c_{ij} + d_{ij}) \phi(\bar{c}_{ij} + d_{ij}) + c.c.] \quad (5)$$

where $c.c.$ is the complex conjugate. Before considering the frustrated case in detail, we briefly review the variational wavefunction for the unfrustrated case [12]. There, $t > 0$, so the matrix elements of H_t are all non-positive in the dimer basis. Thus the Perron-Frobenius theorem applies and the ground state can be taken to be everywhere positive. Hence, we get a normalizable wavefunction if $\phi(C) = \sqrt{P(C)}$ with $P(C)$ taken as the probability of the dimer configuration C . Equivalently, one can assign positive weights $W(C)$ to each dimer configuration C , then the probability $P(C) = W(C)/Z$, where $Z = \sum_C W(C)$.

The central assumption that leads to tractable wavefunctions is the following. We assign edge weights w_{ab} to all honeycomb lattice edges ab , and write the weight of dimer covering C as $W(C) = \prod_{\text{covered } \langle ab \rangle} w_{ab}$. Interpreting W as a fictitious Gibbs weight, this corresponds to a problem of hardcore dimers in an external potential. Powerful Grassmann variable techniques have been developed for this problem, which will allow us to calculate properties of these wavefunctions.

Plug the ansatz $\phi(C) = \sqrt{P(C)}$ into (5), and using the fact that the ratio $\frac{P(\bar{c}_{ij} + d_{ij})}{P(c_{ij} + d_{ij})}$ is independent of the configuration d_{ij} .

$$E = -t \sum_{\langle ij \rangle} \sqrt{P(c_{ij})P(\bar{c}_{ij})} \quad (6)$$

where $P(c_{ij}) = \sum_{d_{ij}} P(c_{ij} + d_{ij})$ is the net probability of the local configuration.

The dimer number operator is $n_{ab} = (1 + s_i^z s_j^z)/2$ where ab is the honeycomb lattice edge dual to the triangular lattice edge ij . The probability $P(c_{ij})$, $P(\bar{c}_{ij})$ are the expectation values $\langle n_{12} n_{34} n_{56} n_{78} n_{9,10} \rangle$, $\langle n_{23} n_{45} n_{67} n_{89} n_{10,1} \rangle$, respectively. This can be evaluated analytically by the Grassmannian integral method [14]. In the Grassmannian formulation, the dimer partition function is represented as an integral over Grassmannian variables η_a defined on the honeycomb lattice sites, $Z = \int \exp(\sum_{a,b} \eta_a A_{ab} \eta_b / 2) \prod_a \eta_a = \text{Pf}[A]$, where $\text{Pf}[A]$ is the Pfaffian of the Kasteleyn matrix A [15], and $A_{ab} = +w_{ab}$ if the Kasteleyn orientation is from a to b , or $= -w_{ab}$ if otherwise (see FIG. 2 (b)). The probability $P(c_{ij})$ is calculated as an expectation value in the Grassmannian theory, the rule is to replace n_{ab} by $A_{ab} \eta_a \eta_b$, then we get $P(c_{ij}) = w_{12} w_{34} w_{56} w_{78} w_{9,10} \left| \langle \prod_{a=1}^{10} \eta_a \rangle \right|$. Thus the variational energy (6) can be written as $E = -t \sum_{\langle ij \rangle} \sqrt{\prod_{i=1}^{10} w_{i,i+1}} \left| \langle \prod_{a=1}^{10} \eta_a \rangle \right|$ with $w_{10,11} = w_{10,1}$. The ten-point correlator of anticommuting η can be Wick-expanded into a Pfaffian of a 10×10 antisymmetric matrix, $\left| \langle \prod_{a=1}^{10} \eta_a \rangle \right| = \text{Pf}[\langle \eta_a \eta_b \rangle] = \sqrt{\det[\langle \eta_a \eta_b \rangle]}$, $a, b =$

$1 \dots 10$. The above formula can be further simplified to the determinant of a 5×5 matrix exploiting the bipartiteness of the honeycomb lattice: $\left| \langle \prod_{a=1}^{10} \eta_a \rangle \right| = |\det[\langle \eta_a \eta_b \rangle]|$, $a = 1, 3, \dots, 9$; $b = 2, 4, \dots, 10$. This is much more efficient than the brutal-force Wick expansion used by Sen *et al.* [12], which allows us to evaluate more complicated correlation functions later in this paper. The two-point correlator $\langle \eta_a \eta_b \rangle = (A^{-1})_{ba}$ can now be evaluated by a Fourier transformation since the Kasteleyn matrix A has 2D translational symmetry. For the chosen Kasteleyn orientation and basis shown in FIG. 2 (b)), in the thermodynamic limit, the two-point correlator of the site a in unit cell $(0, 0)$ and the site b in unit cell (x, y) is

$$\langle \eta_{a,(0,0)} \eta_{b,(x,y)} \rangle = \int_0^{2\pi} \int_0^{2\pi} [\tilde{A}^{-1}(\vec{k})]_{ba} e^{i(k_x x + k_y y)} \frac{dk_x dk_y}{4\pi^2}$$

where $a, b = 1, \dots, 6$, and $\tilde{A}^{-1}(\vec{k})$ is the inverse of the 6×6 anti-hermitian matrix $\tilde{A}(\vec{k})$,

$$\tilde{A}(\vec{k}) = \begin{pmatrix} 0_{3 \times 3} & R(\vec{k}) \\ -R^\dagger(\vec{k}) & 0_{3 \times 3} \end{pmatrix}, \text{ with } R(\vec{k}) = \begin{pmatrix} \frac{1}{\epsilon_x \epsilon_y} & z & z \\ z & \epsilon_y & z \\ z & z & \epsilon_x \end{pmatrix}$$

where $\epsilon_x = e^{ik_x}$, $\epsilon_y = e^{ik_y}$. As is shown in Ref. 12, for $t > 0$ this variational wavefunction has two local minima at $z \approx 0.9258$ with energy per site $E = -0.13774t$, and $z \approx 1.073$ with energy per site $E = -0.13762t$, corresponding to the two supersolid states, $(+ - -)$ and $(0 + -)$ of the triangular lattice boson model [3, 4, 5]. For the frustrated case, the wavefunction is obtained by unitary transformation, $|\Psi'\rangle = U|\Psi\rangle = \sum_C \phi'(C)|C\rangle$, hence the variational wavefunction is $\phi'(C) = i^{N_s(C)} \sqrt{P(C)}$. The variational energy $E = \langle \Psi' | H_t | \Psi' \rangle$ is of course the same as in the unfrustrated case. In order to understand the two variational wavefunctions better, we shall calculate two point correlation functions, assuming for simplicity that the two points i, j are on the same horizontal line, and j is on the right.

Diagonal Correlations: Consider first the density-density correlation function $\langle s_i^z s_j^z \rangle$. Draw a line from i to j and it will cut through an set of honeycomb lattice edges $\langle ab \rangle$. If the number of edges with no dimers cut by this line is even, then $s_i^z s_j^z = +1$ and otherwise $= -1$. In terms of the dimer number operator n_{ab} the $s_i^z s_j^z$ becomes a non-local string operator,

$$\langle s_i^z s_j^z \rangle = \langle \prod_{\langle ab \rangle \text{ cut by } ij} (2n_{ab} - 1) \rangle \quad (7)$$

Expand the product we get $2^{|j-i|}$ terms ($|j-i|$ is the distance between j and i measured by the triangular lattice constant), each of which is the type of correlation functions evaluated before. Because these operators are diagonal in the dimer basis, $\langle \Psi | s_i^z s_j^z | \Psi \rangle = \langle \Psi' | s_i^z s_j^z | \Psi' \rangle$.

Off diagonal Correlations: The square of the off-diagonal long range order (ODLRO) parameter $\langle b_i^\dagger b_j \rangle$ is slightly more complicated. In the dimer basis it describes the simultaneous resonances of two hexagons (if i and j are not neighbors).




FIG. 3: (Color online) Two possible simultaneous double-resonance needed for calculating $\langle b_i^\dagger b_j \rangle$: $C_i, C_j \leftrightarrow \bar{C}_i, \bar{C}_j$, and $C_i, \bar{C}_j \leftrightarrow \bar{C}_i, C_j$, with even(odd) number of no-dimer edges cut by the line $i + \hat{x}, j - \hat{x}$ (dash line).

However for this process to happen s_i^z and s_j^z must be opposite. Label the two local resonating configurations on hexagon $i(j)$ by $C_{i(j)}$ and $\bar{C}_{i(j)}$, there are two possibilities of this simultaneous double-resonance, shown in FIG. 3, with opposite conditions for the edges cut by the line $i + \hat{x}, j - \hat{x}$, where $\hat{x}$ is the horizontal triangular lattice vector. The even(odd) requirement can be enforced using the dimer number operators as $[1 \pm \prod (2n_{ab} - 1)]/2$, where the product is over all edges $< ab >$ cut by the line $i + \hat{x}, j - \hat{x}$ (see FIG. 3 for an example). Consider $t > 0$ case first, we have

$$\begin{aligned} \langle \Psi | b_i^\dagger b_j | \Psi \rangle &= \frac{w_{23}w_{45}w_{61}w_{89}w_{10,11}w_{12,7}}{w_{12}w_{34}w_{56}w_{78}w_{9,10}w_{11,12}} \\ &\times \left\{ \left\langle n_{12}n_{34}n_{56}n_{78}n_{9,10}n_{11,12} \cdot [1 + \prod (2n_{ab} - 1)]/2 \right\rangle \right. \\ &\quad \left. + \left\langle n_{12}n_{34}n_{56}n_{78}n_{9,10}n_{11,12} \cdot [1 - \prod (2n_{ab} - 1)]/2 \right\rangle \right\} \\ &= \sqrt{\prod w} \left| \left\langle \prod_{a=1}^{12} \eta_a \right\rangle \right| \end{aligned} \quad (8)$$

where the $\prod w$ is the product of edge weights of the twelve(12) edges around hexagons i and j . Note, this simple form arises because the string $\prod (2n_{ab} - 1)$ cancels out. We will see shortly that in the frustrated hopping case, this does not happen.

If distance between i and j is large, the 12-point correlator $\langle \prod_{a=1}^{12} \eta_a \rangle$ can be factorized into two 6-point correlators $\langle \prod_{a=1}^6 \eta_a \rangle \cdot \langle \prod_{a=7}^{12} \eta_a \rangle$. And we have the relation $\sqrt{w_{12}w_{34}w_{56}w_{78}w_{9,10}w_{11,12}} |\langle \prod_{a=1}^6 \eta_a \rangle| = \langle \Psi | b_i^\dagger | \Psi \rangle$. So literally we have the factorization property

$$\langle \Psi | b_i^\dagger b_j | \Psi \rangle \rightarrow \langle \Psi | b_i^\dagger | \Psi \rangle \langle \Psi | b_j | \Psi \rangle, \quad |j - i| \rightarrow \infty$$

For $t < 0$ case we need to take care of the phases of ϕ' . From FIG. 2 (c) we can see that $\langle \Psi' | b_i^\dagger b_j | \Psi' \rangle$ has similar form as the first line of (8), only the first term inside $\{ \cdot \}$ acquires a minus sign. Therefore we get

$$\begin{aligned} \langle \Psi' | b_i^\dagger b_j | \Psi' \rangle &= - \frac{w_{23}w_{45}w_{61}w_{89}w_{10,11}w_{12,7}}{w_{12}w_{34}w_{56}w_{78}w_{9,10}w_{11,12}} \\ &\times \left\langle n_{12}n_{34}n_{56}n_{78}n_{9,10}n_{11,12} \prod (2n_{ab} - 1) \right\rangle \end{aligned} \quad (9)$$




FIG. 4: Supersolid LRO from the variational wavefunction in the strong repulsion limit. Greyscale shows density order $\langle 2n_i - 1 \rangle$ while arrows denote superfluid order $\langle b_i^\dagger \rangle$, for (a): Frustrated hopping ($t < 0$) - note the sign structure of superfluid order; and (b): Unfrustrated hopping ($t > 0$)

The product can be expanded into $2^{|j-i|-2}$ terms, each of which can be evaluated as before. Note, this correlation function cannot be factorized as in the unfrustrated case and one necessarily needs to evaluate a string correlator.

Results: We evaluate the above mentioned correlators up to distance $|j - i| = 18$ and extrapolate to infinite distance limit to determine the long range order.

At the global energy minimum $z = 0.9258$ and for the *unfrustrated* ($t > 0$) case, the long range order parameter $\langle \vec{s} \rangle = (b^\dagger + b, ib - ib^\dagger, 2n - 1) = (0.163, 0, 0.764), (0.372, 0, -0.412), (0.372, 0, -0.412)$ for the three sublattices A,B,C, respectively (we have set the superfluid phase to zero and sublattice A is surrounded by weight z hexagon in FIG. 2 (b)). These numbers are in agreement with Quantum Monte Carlo (QMC) results. The average density deviation from $1/2$ is $|0.764 - 0.412 - 0.412|/2/3 = 0.010$, which is about 2%, in good agreement with QMC [4]. The solid order parameter is $|n_A + n_B e^{2\pi i/3} + n_C e^{4\pi i/3}|^2/9 = 0.0384$ ($n_{A,B,C}$ are boson densities on the three sublattices), which is about 15% smaller than the QMC result of 0.045 [4, 10], but in good agreement with classical Monte Carlo calculations result 0.0389 of the same type of variational wavefunctions [12].

In the *frustrated* ($t < 0$) case the three sublattice order is $(0, 0, 0.764), (0.389, 0, -0.412), (-0.389, 0, -0.412)$, as shown in Fig. FIG. 4. The average density deviates from $1/2$ by the same amount as the frustrated case. In spin language this means a non-zero average z-component of spin, $|\sum_i S_i^z/N| = |\sum_i s_i^z/(2N)| = 0.01$, which is about 50% smaller than harmonic spin-wave result 0.02, which has the same symmetry[7]. Note that surprisingly the superfluid amplitude (XY-component of $\vec{s}$) on the B,C sublattices is *larger* than those in the unfrustrated case, while it vanishes on the A sublattice. Note, this quantity can be directly measured in Quantum Monte Carlo simulations of the unfrustrated system in the large repulsion limit, by calculating correlations of the unitarily transformed operator. For example, with $O = s_i^+ s_j^-$, with j to the right of i in the same horizontal line and $|j - i| > 2$, the correlator to be measured is $U^\dagger O U = -s_{i+\hat{x}}^+ s_{j-\hat{x}}^-$. Finally, we combine the present

results in the large repulsion (or $V \gg -t$) limit with known 120° order in the isotropic $V = -2t$ ($J_z = J_\perp$) limit. These can be connected without a phase transition, if we assume that the supersolid phase persists with no change in symmetry over the entire range $V > -2t > 0$. This scenario, which is also a phase diagram for the spin 1/2 XXZ antiferromagnet, is depicted in Figure 1. [18]

Experimental Realization: How can the frustrated boson hoppings be experimentally realized? In lattice cold atom systems, a recent experiment [16] demonstrated that ‘repulsively’ bound molecular bosons have frustrated hoppings. Consider preparing an initial state composed of molecules of pairs of atoms (either bosons or fermions) with one or zero molecules per site. If the interactions between atoms are now made repulsive, the effective molecular hopping is readily seen to be frustrated, since the singly occupied sites are lower in energy. If this metastable state is sufficiently long lived, the equilibrium properties of the frustrated boson system can be accessed. In Josephson Junction Arrays, external magnetic fields can generate frustrated hopping[17].

We acknowledge funding from nsf-dmr 0645691 and ARO Grant No. W911NF-07-1-0576, and useful discussions with Ehud Altman and Arnab Sen.

- [2] E. Kim and M.H.W. Chan, Nature (London) **427**, 225 (2004); Science **305**, 1941 (2004)
- [3] S. Wessel and M. Troyer, Phys. Rev. Lett. **95**, 127205 (2005)
- [4] D. Heidarian and K. Damle, Phys. Rev. Lett. **95** 127206 (2005)
- [5] R. G. Melko, A. Paramekanti, A. A. Burkov, A. Vishwanath, D. N. Sheng, L. Balents, Phys. Rev. Lett. **95**, 127207 (2005)
- [6] P. Fazekas, P.W. Anderson, Philos. Mag. **30**, 423 (1974)
- [7] B. Kleine, E. Müller-Hartmann, K. Frahm, and P. Fazekas, Z. Phys. B - Condensed Matter **87**, 103 (1993)
- [8] N. Read and S. Sachdev, Phys. Rev. Lett. **62**, 1694 (1989); N. Read and S. Sachdev, Phys. Rev. B **42**, 4568 (1990).
- [9] E. Fradkin and S. A. Kivelson, Mod. Phys. Lett. B **4**, 225 (1990).
- [10] M. Boninsegni and N. Prokof'ev, Phys. Rev. Lett. **95** 237204 (2005)
- [11] D.N. Sheng et. al., unpublished.
- [12] A. Sen, P. Dutt, K. Damle, R. Moessner, Phys. Rev. Lett. **100**, 107204 (2008)
- [13] G. H. Wannier, Phys. Rev. **79**, 357 (1950)
- [14] S. Samuel, J. Math. Phys. **21**, 2806 (1980)
- [15] P.W. Kasteleyn, J. Math. Phys. **4**, 287 (1963)
- [16] K. Winkler et. al., Nature **441**, 853-856 (2006)
- [17] B. Pannetier, J. Chaussy, R. Rammal, and J. C. Villegier, Phys. Rev. Lett. **53**, 1845 (1984).
- [18] At the other local minimum $z = 1.073$, the three sub-lattice order for unfrustrated and frustrated cases are $(0.438, 0, 0)$, $(0.231, 0, 0.661)$, $(0.231, 0, -0.661)$ and $(0.340, 0, 0)$, $(-0.147, 0, 0.661)$, $(-0.147, 0, -0.661)$.

[1] A.F. Andreev and I.M. Lifshitz, Sov. Phys JETP **29**, 1107 (1969)